

Volatile organic compounds from *Pachyrhizus ferrugineus* and *Pachyrhizus erosus* (Fabaceae) leaves

[Compuestos orgánicos volátiles de las hojas de *Pachyrhizus ferrugineus* y *Pachyrhizus erosus* (Fabaceae)]

Edgar A. Estrella-Parra¹, Juan C. Gómez-Verjan¹, Carlos L. Céspedes², Julio Alarcón², Zenón Cano-Santana³, Alberto Reyes-García⁴, Marco A. Cerbón-Cervantes⁵ & Ricardo Reyes-Chilpa¹

¹Instituto de Química. Universidad Nacional Autónoma de México. México

²Departamento de Ciencias Básicas, Facultad de Ciencias, Universidad del Bío-Bío, Chillán, Chile

³Departamento de Ecología y Recursos Naturales, Facultad de Ciencias. Universidad Nacional Autónoma de México. México

⁴Instituto de Biología, Herbario MEXU, Universidad Nacional Autónoma de México. México

⁵Facultad de Química. Universidad Nacional Autónoma de México. México

Contactos / Contacts: Ricardo REYES-CHILPA - E-mail address: chilpa@unam.mx

Abstract: In México, *Pachyrhizus erosus* (Fabaceae) commonly called "jícama", is widely known for its edible tubers. It is cultivated since the pre-Columbian period, and the powdered seeds have been used for the treatment of mange, lice, and fleas, due to their content of rotenone, a well-known insecticidal compound. On the other hand, *P. ferrugineus*, a wild species can only be found in the Tropical Forests, and has no commercial value. It is known that plants release volatile organic compounds (VOCs) showing qualitative and quantitative differences if are wild or cultivated. VOCs are also involved as repelling or attracting chemical signals to insect herbivores, and their natural enemies. Until now, the VOCs of the leaves of *P. erosus* and *P. ferrugineus* have not been investigated. In the present contribution the VOCs of both species were characterized by headspace solid-phase (HS-SPME) extraction and gas chromatography-mass spectrometry (GC-MS-TOF). In *P. erosus* 21 VOCs were found, being the most abundant: cyclohexanone (32.8%), 3-hexen-1-ol (Z) (32.7%), 3-hexenal (Z) (10.5%). The majoritarian compounds were C6 or C5 derivatives. In *P. ferrugineus*, the most abundant VOCs were: 5-hexene-1-ol acetate (51.5%), undecanal (22.4%), 2-hepten-1-ol (14.5%). The majoritarian compounds were C6, C7 or C11 derivatives.

Keywords: *Pachyrhizus ferrugineus* leaves, *Pachyrhizus erosus* leaves, volatile organic compounds.

Resumen: En México, *Pachyrhizus erosus* (Fabaceae) es llamada comúnmente "jícama" y es conocida por sus tubérculos comestibles. Se ha cultivado desde el período pre-Colombino y las semillas se han utilizado para el tratamiento tópico de la sarna, piojos, pulgas; las semillas contienen rotenona, un compuesto insecticida. Por otra parte, *P. ferrugineus* solo está presente en estado silvestre en los bosques tropicales y carece de valor comercial. Se sabe que las plantas liberan compuestos orgánicos volátiles (COV) y muestran diferencias cualitativas y cuantitativas dependiendo, si son silvestres o cultivadas. Los COV también son señales químicas atrayentes o repelentes de los insectos herbívoros y a sus enemigos naturales. Hasta ahora, los COV en las hojas de *P. erosus* y *P. ferrugineus* no han sido investigados. En el presente trabajo, los COV se identificaron mediante la microextracción (HS-SPME) en fase sólida, e identificados por cromatografía de gases acoplada a espectrometría de masas (GC-MS-TOF). En *P. erosus* se encontraron 21 COV, siendo los más abundantes: ciclohexanona (32.8%), 3-hexen-1-ol (Z) (32.7%) y 3-hexenal (Z) (10.5%). Los compuestos mayoritarios son C6 y C5. En *P. ferrugineus* los más abundantes fueron: 5-hexen-1-ol acetato (51.5%), undecanal (22.4%) y 2-hepten-1-ol (14.5%). Los compuestos mayoritarios son C6, C7 o C11.

Palabras clave: Hojas de *Pachyrhizus ferrugineus*, Hojas de *Pachyrhizus erosus*, compuestos volátiles

Recibido | Received: November 28, 2014

Aceptado | Accepted: August 16, 2015

Aceptado en versión corregida | Accepted in revised form: February 4, 2016

Publicado en línea | Published online: May 30, 2016.

Este artículo puede ser citado como / This article must be cited as: EA Estrella-Parra, JC Gómez-Verjan, CL Céspedes, J Alarcón, Z Cano-Santana, A Reyes-García, MA Cerbón-Cervantes, R Reyes-Chilpa. 2016. Volatile organic compounds from *Pachyrhizus ferrugineus* and *Pachyrhizus erosus* (Fabaceae) leaves. *Bol Latinoam Caribe Plant Med Aromat* 15 (3): 175 – 181.

INTRODUCTION

The Neotropical genus *Pachyrhizus* is placed taxonomically in the subtribe Diocleinae, tribe Phaseoleae, within the legume family (Fabaceae) (Sørensen *et al.*, 1996). The genus comprises five species found in the American Continent; three of these are cultivated to obtain edible tuberous roots, and the other two only can be found wild. *Pachyrhizus erosus* and *P. ferrugineus* are present in Central America and Mexico, while *P. tuberosus*, *P. ahipa* and *P. panamensis* are distributed in South America (Sørensen *et al.*, 1996; Ramos *et al.*, 2013).

In México, there are only two *Pachyrhizus* species: *P. erosus* (cultivated) and *P. ferrugineus* (wild) (Sørensen *et al.*, 1996). The first one is commonly called "jícama", it is widely known for its edible tubers. There is evidence that during the pre-Columbian period it was cultivated by the Toltec, Aztec and Maya civilizations (Sørensen *et al.*, 1996). Today, *P. erosus* is a tuber legume crop with high yield potential, high nutritional value, low N₂ fertilizer and low pesticide demand (Castellanos *et al.*, 1997), being ideal for plant growing. The seeds of *P. erosus* have been used in Mexico for the treatment of mange since pre-Columbian times (Bejar *et al.*, 2000), and during the XX century the peasants used the powdered seeds for their insecticidal properties against ectoparasites, such as lice, and fleas. These ethnomedical applications can be explained due to their content of rotenone, a compound well known compound for its insecticidal and acaricidal properties (Reyes-Chilpa *et al.*, 2003), as well as other isoflavonoids.

Pachyrhizus erosus is extensively cultivated in Mexico, even for exportation, and has achieved wide acceptance in U.S.A., Southeast Asia, and Western Africa; therefore, it has stimulated agronomic and food science interest (Ramos *et al.*, 2013). It has been estimated that México exported 11,000 tons per year to USA, where it reaches an average price of one dollar per kg (Castañeda, 2000).

Volatile organic compounds (VOCs) are lipophilic compounds with low molecular weight and high vapor pressure at ambient temperatures (Dudareva *et al.*, 2013). The VOCs are emitted as blends from flowers, leaves, fruits, and roots into the surrounding atmosphere (Kigathi *et al.*, 2009). Artificial selection for specific plant traits has often not only changed the appearance of cultivated plants compared to their wild conspecifics, but also their

chemistry (Gols *et al.*, 2011). Furthermore, volatile organic compounds have been widely studied due to their contribution to aroma and flavour and as markers for authenticity, for example, biological or geographical origin, cultivar (Socaci *et al.*, 2013). To our best knowledge, volatile constituents of leaves of *P. erosus* (cultivated type) and *P. ferrugineus* (wild type) has not been reported. In the present study volatile compounds of these species were analyzed by HS-SPME/GC-MS-TOF as a starting point of future studies aimed to study their susceptibility to insects, and fungi.

MATERIALS AND METHODS

Plant material

The leaves of two specimens were free of any pesticide. *Pachyrhizus erosus* leaves were collected from farmland to San Andres Tlalquitenango, Morelos, México. A voucher specimen was deposited in MEXU-UNAM (138292). At the time of collecting the plants were in bloom, the flowers were of purple color. *Pachyrhizus ferrugineus* leaves were collected in a Deciduous Tropical Forest in the Natural Reserve "La Sepultura", Villaflores, Chiapas, México. A Voucher was deposited in MEXU-UNAM (1060890). The plants were collected in the core area of the reserve (wild plants). At the time of collecting the plants had mature fruits, which the contained of 6-8 seeds. The specimens were identified and authenticated by one of us, Alberto Reyes-Garcia from the Instituto de Biología, Universidad Nacional Autónoma de México (UNAM, México).

Collection of VOCs blend by HS-SPME

To identify and quantify the volatile compounds (VOCs) in *P. erosus* and *P. ferrugineus* leaves, the headspace HS-SPME technique was employed. Volatiles were collected twice ($N = 2$). For this study, the temperature was maintained at 18 ± 1 °C. The VOCs slurry was prepared by blending freshly leaves (5.0 g), 20% sodium chloride solution (2 mL) in a 20 mL flask with a cap and Teflon-faced silicone rubber septa (Supelco, Co., Bellefonte, PA). The flask was placed on a magnetic stirring plate and stirred at 1100 rpm for 15 m. A SPME fiber (DVB/CARBOXEN/PDMS; Supelco, Co., Bellefonte, PA) was then exposed to the headspace of the slurry for 15 m in a water bath at 45° C. The analytical methodology here used was selected after

testing 4 different fibers and previous experiences for VOCs analysis (Torres-González et al., 2015).

Leaf volatiles organic compounds analysis by CG-MS-TOF

The analysis of the volatile compounds (VOCs) of *P. erosus* and *P. ferrugineus* leaves was performed with a gas chromatography coupled to a mass spectrometer (GC-MS). The sample analysis was conducted using a Gas Chromatograph (Agilent 6890N) with helium (grade 5; ultra-high purity) as the carrier gas; the outlet of the column (DB-5, 20 m×0.18 mm×0.18 µm film) was coupled to a Mass Spectrometer (LECO Pegasus 4D). Parameters for electron impact sample ionization were as follows. Ionization chamber temperature: 200° C; interface temperature: 250° C; source temperature: 230° C. The gas chromatographic conditions were as follows: split less injection (1:20) at 300° C; initial oven temperature, 40° C for 1 min, increased at 8° C per minute to 300° C and held for 5 min. Desorption time was 4 min. Mass Analyzer: Time of Flight (TOF); Spectral Acquisition: 20 spectra/second; delayed ignition of the filament: 0 min; mass range: 33-400; Ionization chamber temperature: 200° C; calibration compound: perfluoroterbutylamine (PFTBA). To

identify the compounds their mass spectra were compared with NIST/EPA/NIH, and reported Kovats Retention Index (KI, www.pherobase.org and www.flavornet) libraries. The composition was reported as relative percentage of the total peak area. Individual compounds were quantified by calculating the peak area relative to the peak area for the internal standard. A mixture of pure alkanes standards C8 to C24 were used for determining the Kovats index.

RESULTS

In *P. erosus* 21 kind of VOCs were found, being the most abundant: cyclohexanone (32.8%), 3-hexen-1-ol (Z) (32.7%), 3-hexenal (Z) (10.5%), furan-2-ethyl (6.4%), and pentanol (6.4%). Together are 88.99% of the VOCs blend. The majoritarian compounds were C6 or C5 derivatives including ketones, aldehydes, and alcohols (Table 1). Eleven kind of VOCs were detected from *P. ferrugineus*, being the most abundant: 5-hexene-1-ol acetate (51.5%), undecanal (22.4%), 2-hepten-1-al (14.5%), and 2-eicosene (Z) (4.76). Together represent 93.24% of the VOCs blend. The majoritarian compounds were C6, C7 or C-11 derivatives including ester, and aldehydes (Table 1).

Tabla 1
Volatile organic compounds (VOCs) contained in *P. erosus* (cultivated plant) and *P. ferrugineus* (wild plant) leaves.

<i>Pachyrhizus erosus</i>						<i>Pachyrhizus ferrugineus</i>				
Peak	RT (min)	Compound	KIc	KIr	Abun (%)	RT (min)	Compound	KIc	KIr	Abun (%)
1	0.12	Pentanal	733.9	732	1.66	4.11	2-hepten-1-al	946.5	951	14.57
2	0.33	3-methyl-1-butanol	736.0	736	1.66	5.73	5-Hexene-1-ol, acetate	1092.1	1070	51.50
3	0.58	Pentanol	759.4	759	6.43	7.56	Undecanal	1317.6	1310	22.41
4	1.01	Furan 2-ethyl	761.7	730	6.44	8.19	2-dodecanone	1406.3	1401	0.06
5	2.01	3-hexenal, (Z)-	819.1	800	10.57	9.13	2H-Pyran-2-one, tetrahydro-6-pentyl-1-	1558.1	1525	1.24
6	3.44	Ciclohexanone	901.0	896	32.84	9.99	Heptadecene	1709.1	1696	0.95
7	3.47	3-Hexen-1-ol, (Z)-	903.1	882	32.71	10.54	4,14-dimethylheptadecane	1816.4	1816	0.87
8	3.589	(E,E)-2,4-Hexadienal	910.4	910	0.07	11.22	n-Hexadecanoic acid	1953.3	1951	0.80

9	3.91	Heptanal	932.8	912	0.23	11.57	2-eicosene (Z)	2023.8	2022	4.76	
10	4.18	2 heptenal	951.2	951	3.84	11.59	5-Ethylnonadecane	2027.7	2034	0.73	
11	4.99	(E,E)-2-4 heptadienal	1009.5	1009	1.96	12.20	Methyl stearate	2161.5	2139	2.11	
12	5.21	(E)-3-octen-2-one	1034.1	1034	0.51						
13	5.41	(E)-2-octenal	1056.5	1056	0.04						
14	5.56	1-octanol	1072.7	1072	0.06						
15	6.2	Hexanoic acid, 2-methylpropyl ester	1149.3	1149	0.18						
16	7.14	2-Cyclohexen-1-one, 3-methyl-6-(1-methylethyl)-	1259.9	1258	0.08						
17	8.09	2-Methylpropanoic acid 3-hydroxy-2,4,4-trimethylpentyl ester	1390.7	1380	0.20						
18	8.25	Unknown	1416.5	ND	0.28						
19	8.57	(E)- β -Ionone sesquiterpene	1467.0	1467	0.06						
20	8.93	3,5,9-Undecatrien-2-one, 6,10-dimethyl-terpene	1526.0	1527	0.10						
21	9.53	2-Methylpropanoic acid, 1-(1,1-dimethylethyl)-2-methyl-1,3-propanediyl ester	1626.4	1594	0.08						
		Total area (%)						Total area (%)			
		Identified total area (%)						Identified total area (%)			
		Identified compounds (%)						Identified compounds (%)			
			100						100		
			99.72						100		
			95.23						100		

KIc: Calculated Kovats retention index;
KIr: Reported Kovats retention Index;
ND: Not determined. RT: retention time

DISCUSSION

It is well known that plants release different VOCs blends according if they belong to the wild or cultivated types (Gols *et al.*, 2011); as well as, if they are attacked or not by herbivores (Kigathi *et al.*, 2009) or are under drought stress Copolovici *et al.*, 2014). Several authors have reported losses of VOCs in cultivated as compared to wild ancestors, for instance in the case of cranberry (*Vaccinium macrocarpon*), caterpillars performed best on the highest-yielding variety which has low induction of volatile sesquiterpenes; therefore it has proposed that breeding in cranberry has compromised plant defenses (Rodríguez *et al.*, 2011). Gols *et al.* (2011) have reported not only differences among wild and cultivated cabbages (*Brassica oleracea*), but also that the parasitoid *Cotesia rubecula* and specialized on hosts feeding on brassicaceous plants, was differentially attracted and preferred wild types over cultivated cabbage. VOCs are emitted as a response to insect egg deposition and it also been postulated that their content may drastically diminish as a consequence of crop breeding in maize (Tamiru *et al.*, 2011). It has also been reported that communication among plants via VOCs is a defense strategy to herbivores (Muroi *et al.*, 2011); thus, plants can reduce the emission of VOCs blend when growing together as an strategy to reduce herbivory attack (Kigathi *et al.*, 2013).

Differences between *P. erosus* and *P. ferrugineus* VOCs blend, may be due, in first instance that are different species, and second term that are cultivated and wild plants, respectively. It is interesting that *P. erosus* (cultivated) leaves release more VOCs (21) than the wild species *P. ferrugineus* (11). Regarding to the type of VOCs in the species of *Pachyrhizus* here studied, both contain mainly aldehydes, ketones, and alcohols. Only *P. erosus* leaves contained terpenoids such as (E)- β -ionone. Several compounds, such as 3-hexenal (Z), undecanal and 3-hexen-1-ol (Z) have been reported as common constituents of many other plant species (Srinivasa *et al.*, 1989; Kulkarni *et al.*, 1998; Kaul *et al.*, 1999; Asuming *et al.*, 2005; Rohloff & Bones, 2005; Tayoub *et al.*, 2006). 3-Hexen-1-ol is part of the

chemoattractant blend of VOCs to insect natural enemies, when certain plants are invaded or ovoposited by herbivores (Warthen *et al.*, 1997; Han & Chen, 2002; Mumm *et al.*, 2004). Some of the compounds found here in low amounts have been reported to possess insecticidal and antifeedant activity; for instance (E)- β -Ionone Gruber *et al.*, 2009; (Alarcón *et al.*, 2013; Céspedes *et al.*, 2013; Muñoz *et al.*, 2013; Céspedes *et al.*, 2014). Cyclohexanone have only found to date in mango cultivars (Pino *et al.*, 2005).

The present research provides the first approach to the study of the role of VOCs blends in *P. erosus* and *P. ferrugineus*. Until now, it has only been described that *P. erosus* seeds contains isoflavonoids with insecticidal activity (Bejar *et al.*, 2000), for instance rotenone (Alavez *et al.*, 1998, Estrella-Parra *et al.*, 2014); however it is unknown if rotenone and other insecticidal isoflavonoids are also synthesized in *P. erosus*, and *P. ferrugineus* leaves.

CONCLUSIONS

This is the first study about the composition of VOCs from *P. erosus* and *P. ferrugineus* leaves and improves the knowledge of the genus *Pachyrhizus*. The VOCs blend from the wild (*P. ferrugineus*) and cultivated plant species (*P. erosus*) showed different composition. *Pachyrhizus erosus* leaves released more VOCs than *P. ferrugineus* leaves. Both species contain ketones, aldehydes, alcohols, and esters.

ACKNOWLEDGEMENTS

EA Estrella-Parra is grateful to Posgrado en Ciencias Biológicas de la Universidad Nacional Autónoma de México (PCB-UNAM), and Consejo Nacional de Ciencia y Tecnología (CONACYT; 220355) for providing a scholarship. We thank the funding provided by the Dirección de Investigación of the University of the Bío Bío (Grupo de Investigación Química y Biotecnología de Productos Naturales Bioactivos) FONDECYT Projects Grants # 1130463 and 1130242. We thank to Georgina Artemisa Duarte Lisci (USAI-UNAM) for technical assistance.

REFERENCES

Alarcón J, Lamilla C, Céspedes C. 2013. Insecticidal activity of sesquiterpenes skeleton synthesized by the conventional Robinson annulations reaction on *Drosophila*

- melanogaster*. **Ind Crops Prod** 42: 268 - 272.
- Alavez-Solano D, Reyes-Chilpa R, Jiménez M, Villavicencio M, Pérez E, Gómez F. 1998. Estudio químico y actividad insecticida de semillas de *Pachyrhizus erosus* (L.) Urban. **Proceedings of the 2nd International Symposium on Tuberous Legumes**. MacKeenzie, Celaya, Guanajuato, México.
- Asuming WA, Beauchamp PS, Descalzo, JT, Dev BC, Dev V, Frost S, Ma CW. 2005. Essential oil composition of four *Lomatium* Raf. species and their chemotaxonomy. **Biochem Syst Ecol** 33: 17 - 26.
- Bejar E, Reyes-Chilpa R, and Jiménez Estrada M. 2000. **Bioactive Compounds from Selected Plants used in XVI Century Mexican Traditional Medicine**. In: Studies in Natural Products Chemistry. Bioactive Natural Products. Atta-ur-Rahman Ed., Elsevier Science Publishers, Amsterdam, The Netherland.
- Castañeda E. 2000. **Utilización de la jícama (*Pachyrhizus erosus* L. Urban) en la elaboración de pan blanco para favorecer su calidad nutricia**. Tesis de Maestría en Ciencias con especialidad en alimentos. Universidad Autónoma de Nuevo León, Monterrey, Nuevo León, México.
- Castellanos JZ, Zapata F, Badillo V, Pena-Cabriaes JJ, Jensen ES, Heredia-García E. 1997. Symbiotic nitrogen fixation and yield of *Pachyrhizus erosus* L. Urban cultivars and *Pachyrhizus ahipa* Wedd Parodi landraces as affected by flower pruning. **Soil Biology and Biochemistry** 29: 973 - 981.
- Céspedes C, Molina S, Muñoz E, Lamilla C, Alarcón J, Palacios S, Carpinella M, Ávila JG. 2013. The insecticidal, molting disruption and insect growth inhibitory activity of extracts from *Condalia microphylla* Cav. (Rhamnaceae). **Ind Crops Prod** 42: 78 - 86.
- Céspedes C, Salazar J, Ariza-Castolo A, Yamaguchi L, Ávila JG, Aqueveque P, Kubo I, Alarcón J. 2014. Biopesticides from plants: *Calceolaria integrifolia*. **Environ Res** 132: 391 - 406.
- Copolovici L, Kännaste A, Remmel T, Niinemets T. 2014. Volatile organic compound emissions from *Alnus glutinosa* under interacting drought and herbivory stresses. **Environ Exp Bot** 100: 55 - 63.
- Dudareva N, Antje K, Joelle KM, Kaplan I. 2013. Biosynthesis, function and metabolic engineering of plant volatile organic compounds. **New Phytologist** 198: 16 - 32.
- Estrella-Parra EA, Gómez-Verjan JC, González-Sánchez I, Vázquez-Martínez ER, Vergara-Castañeda E, Cerbón MA, Alavez-Solano D, Reyes-Chilpa R. 2014. Rotenone isolated from *Pachyrhizus erosus* displays cytotoxicity and genotoxicity in K562 cells. **Nat Prod Res** 28: 1780 - 1785.
- Gols R, Bullock JM, Dicke M, Bukovinszky T, Harvey JA. 2011. Smelling the wood from the Trees: non-linear parasitoid responses to volatile attractants produced by wild and cultivated cabbage. **J Chem Ecol** 37: 795 - 807.
- Gruber MY, Xu N, Grenkow L, Li X, Onyilagha J, Soroka JJ, Westcott ND, Hegedus DD. 2009. Responses of the crucifer flea beetle to *Brassica* volatiles in an olfactometer. **Environ Entomol** 38: 1467 - 1479.
- Han BY, Chen ZM. 2002. Composition of the volatiles from intact and mechanically pierced tea aphid-tea shoot complexes and their attraction to natural enemies of the tea aphid. **J Agric Food Chem** 50: 2571 - 2575.
- Kaul VK, Gujral RK, Singh B. 1999. Volatile constituents of the essential oil of flowers of *Rosa brunonii* Lindl. **Flavour Fragr J** 14: 9 - 11.
- Kigathi RN, Unsicker SB, Reichelt M, Kesselmeier J, Gershenzon J, Weisser WW. 2009. Emission of Volatile Organic Compounds after herbivory from *Trifolium pratense* (L.) under laboratory and field conditions. **J Chem Ecol** 35: 1335 - 1348.
- Kigathi RN, Weisser WW, Veit D, Gershenzon J, Unsicker SB. 2013. Plants suppress their emission of volatiles when growing with conspecifics. **J Chem Ecol** 39: 537 - 545.
- Kulkarni RN, Mallavarapu GR, Baskaran K, Ramesch S, Kumar S. 1998. Composition of the essential oils of two isomenthone-rich variants of geranium (*Pelargonium* sp.). **Flavour Fragr J** 13: 389 - 392.

- Mumm R, Tiemann T, Schulz S, Hilker M. 2004. Analysis of volatiles from black pine (*Pinus nigra*): significance of wounding and egg deposition by a herbivorous sawfly. **Phytochemistry** 65: 3221 - 3230.
- Muñoz E, Escalona D, Salazar J., Alarcon J, Céspedes C. 2013. Insect growth regulatory effects by diterpenes from *Calceolaria talcana* Grau & Ehrhart (*Calceolariaceae*: *Scrophulariaceae*) against *Spodoptera frugiperda* and *Drosophila melanogaster*. **Ind Crops Prod** 45: 283 - 292.
- Muroi A, Ramadan A, Nishihara M, Yamamoto M, Ozawa R, Takabayashi J, Arimura G. 2011. The composite effect of transgenic plant volatiles for acquired immunity to herbivory caused by inter-plant communications. **PLoS One** 6(10): e24594
- Pino JA, Mesa J, Muñoz Y, Martí MP, Marbot R. 2005. Volatile components from mango (*Mangifera indica* L.) cultivars. **J Agric Food Chem** 53: 2213 - 2223.
- Ramos AM, Renard C, Wicker L, Contreras EJ. 2013. Advances and perspectives of *Pachyrhizus* spp. in food science and biotechnology. **Trends Food Sci Technol** 29: 44 - 54.
- Reyes-Chilpa R, Jiménez Estrada M, Cristóbal Telésforo E, Torres-Colín L, Villavicencio MA, Pérez-Escandón BA, Mercado R. 2003. Natural Insecticides from *Hippocratea excelsa* and *Hippocratea celastroides*. **Econ Bot** 57: 54 - 64.
- Rodríguez C, Vorsa N, Singh AP, Johnson J, Szendrei Z, Mescher MC, Frost CJ. 2011. Tracing the history of plant traits under domestication in cranberries: potential consequences on anti-herbivore defenses. **J Exp Bot** 62: 2633 - 2644.
- Rohloff J, Bones A.M. 2005. Volatile profiling of *Arabidopsis thaliana* putative olfactory compounds in plant communication. **Phytochemistry** 66: 1941 - 1955.
- Socaci SC, Tofana M, Rați IV, Pinteia A. 2013. In-tube extraction and GC-MS analysis of volatile components from wild and cultivated sea buckthorn (*Hippophae rhamnoides* L. ssp. *Carpatica*) berry varieties and juice. **Phytochem Anal** 24: 319 - 328.
- Sørensen M. 1996. **Yam bean (Pachyrhizus DC.). Promoting the conservation and use of underutilized and neglected crops.** 2. Institute of Plant Genetics and Crop Plant Research, Gatersleben/International Plant Genetic Resources Institute, Rome, Italy.
- Srinivasa A, Rajanikanth B, Seshadri R. 1989. Volatile Aroma Components of *Curcuma amada* Roxb. **J Agric Food Chem** 37: 740 - 743.
- Tamiru A, Bruce TJA, Woodcock CM, Caulfield JC, Midega CAO, Ogol CKPO, Mayon P, Birkett MA, Pickett JA, Khan ZR. 2011. Maize landraces recruit egg and larval parasitoids in response to egg deposition by a herbivore. **Ecol Lett** 14: 1075 - 1083.
- Tayoub G, Schwob I, Bessière J, Rabier J, Masotti V, Girard G, Viano J. 2006. Essential oil composition of leaf, flower and stem of styrax (*Styrax officinalis* L.) from South-Eastern France. **Flavour Fragr J** 21: 809 - 812.
- Torres-González A, Lopez-Rivera P, Duarte-Lisci G, López-Ramírez A, Correa-Benítez A, Rivero-Cruz JF. 2015. Analysis of volatile components from *Melipona beecheii* geoprois from Southeast Mexico by headspace solid-phase microextraction. **Nat Prod Res** 29: 1 - 4.
- Warthen JD, Lee CJ, Jang EB, Lance DR, McInnis DO. 1997. Volatile, potential attractants from ripe coffee fruit for female Mediterranean fruit fly. **J Chem Ecol** 23: 1891 - 1900.