



Revisión | Review

## Phytochemistry and Biological Activities of *Werneria* and *Xenophyllum* species

[Fitoquímica y actividades biológicas de especies de *Werneria* y *Xenophyllum*]

Olga Lock<sup>1</sup> & Rosario Rojas<sup>2</sup>

<sup>1</sup>Departamento de Química, Pontificia Universidad Católica del Perú, Lima, Perú

<sup>2</sup>Unidad de Investigación en Productos Naturales, Laboratorios de Investigación y Desarrollo, Facultad de Ciencias y Filosofía, Universidad Peruana Cayetano Heredia, Lima, Perú

Contactos | Contacts: Olga LOCK - E-mail address: [olock@pucep.edu.pe](mailto:olock@pucep.edu.pe)

Contactos | Contacts: Rosario ROJAS - E-mail address: [rosario.rojas@upch.pe](mailto:rosario.rojas@upch.pe)

**Abstract:** Plants of the genera *Werneria* (Asteraceae) and *Xenophyllum* (genus extracted from *Werneria*) are used in traditional medicine of Latin America for the treatment of mountain sickness, hypertension and gastrointestinal disorders. Only a small number of species of these genera have been studied, leading to the isolation of compounds belonging to the classes of benzofurans, chromenes, acetophenones, coumarates, diterpenes and pyrrolizidine alkaloids. Some of the plant extracts and/or compounds have shown antimicrobial, anti-HIV, hypotensive and photoprotective activities.

**Keywords:** *Werneria*; *Xenophyllum*; Medicinal plants; Mountain sickness; Hypertension.

**Resumen:** Las plantas de los géneros *Werneria* (Asteraceae) y *Xenophyllum* (género extraído de *Werneria*) son usadas en la medicina tradicional de América Latina para el tratamiento del mal de montaña, hipertensión y desórdenes gastrointestinales. Solo un pequeño número de especies de estos géneros ha sido investigado, lográndose aislar compuestos que pertenecen a las clases de benzofuranos, cromenos, acetofenonas, cumaratos, diterpenos y alcaloides pirrolizidínicos. Algunos de los extractos y/o compuestos de dichas plantas han mostrado actividades antimicrobianas, anti-HIV, hipotensoras y fotoprotectoras.

**Palabras clave:** *Werneria*; *Xenophyllum*; Plantas medicinales; Mal de montaña; Hipertensión.

Recibido | Received: March 6, 2019

Aceptado | Accepted: April 6, 2019

Aceptado en versión corregida | Accepted in revised form: April 16, 2019.

Publicado en línea | Published online: May 30, 2019

Este artículo puede ser citado como / This article must be cited as: O Lock, R Rojas. 2019 Phytochemistry and biological activities of *Werneria* and *Xenophyllum* species. *Bol Latinoam Caribe Plant Med Aromat* 18 (3): 223 – 238. <http://doi.org/10.37360/blacpma.19.18.3.15>

## INTRODUCTION

The genus *Werneria* was established by Carl S. Kunth and dedicated to the German geologist Abraham G. Werner. The morphological characteristics of this genus show the variability of the species and support that it is not a monophyletic genus, for this reason, almost half of the species were transferred to the new genera *Xenophyllum*, *Misbrookea* and *Anticona* (Funk 1997a; Funk, 1997b; Linares-Perea et al., 2014; Beltrán, 2017). Currently, the *Werneria* genus could be composed of 25-30 species, distributed in the South American Andes (2.800 m.a.s.l) from Venezuela, Colombia, Ecuador, Perú, Bolivia to northern Argentina and Chile, with an isolated population of *W. nubigena* in Mexico and northern Guatemala. The Peruvian territory hosts the greatest wealth of *Werneria* genus with 22 species, of which 5 are endemic (Beltrán, 2017). Recently, *Werneria microphylla* was described as new species from the high Andes of Peru (Beltrán & Leiva, 2018), it is likely that there are more *Werneria* species to be discovered.

The genus *Xenophyllum* consists of 22 species extracted from *Werneria*, 14 of them have been reported in Peru (Beltrán, 2016). “Xeno” means “strange” or “foreign”, while “phyllum” means “leaves”. Members of this genus grow at high elevations (3,000-5,200 m) in the Andes from Colombia to northern Argentina and Chile (Funk, 1997a).

*Misbrookea* is a new genus extracted from *Werneria*, named after its collector Winifred M.A. Brooke. For this genus, only one species (*Misbrookea strigosissima* (A. Gray) V.A. Funk) has been reported; it grows at the grasslands of the highlands of southern Peru and Bolivia (Funk 1997b; Linares-Perea et al., 2014). More recently, Linares-Perea et al. collected *Werneria glareophila* described by Cuatrecasas and found that its characters deviate from the genera *Werneria*, *Misbrookea*, and *Xenophyllum*. They described the new genus *Anticona* comprised by only one species, named as *Anticona glareophila* (Cuatrec.) E. Linares, J. Campos & A. Galán (Linares-Perea et al., 2014).

So far, the phytochemical study and/or biological activities of 9 *Werneria* species have been reported: *W. ciliolate* A. Gray (5 publications), *W. poposa* Philippi (4), *W. nubigena* H.B.K. (3), *W. dactylophylla* Schultz-Bip (3), *W. stuebelli* Hieron (1), *W. decora* cf. Blake (1), *W. staffordiae* Sandwith (1), *W. digitata* Wedell (1) and *W. pigmeae* Gillies ex

Hooker & Arnott (1). In the case of the genus *Xenophyllum*, only two species have been investigated: *X. poposum* (Phillipi) V.A. Funk (7 publications) and *X. incisum* (Phillipi) V.A. Funk (1). It has to be noted that *W. poposa* is now known as *X. poposum*. Of the 28 publications found in the scientific literature for *Werneria* and *Xenophyllum* species, 16 refer to samples collected in Peru, 7 in Argentina, 3 in Chile, 1 in Bolivia and 1 in Ecuador.

Phytochemical studies on the genus *Werneria* have reported a variety of chemical constituents that include pyrrolizidine-type alkaloids, phenolics (such as benzofurans, benzopyrans, *p*-coumarates, coumarins, flavonoids, *p*-hydroxyacetylphenones, caffeoylquinic acids), as well as volatile terpenes, diterpenes (mainly kauran and manoyl derivatives), triterpenes and steroids; all these compounds will be presented in the following sections. Some of them have been reported in a previous review (Lock Sing, 2006).

## CHEMICAL CONSTITUENTS

### Alkaloids

A total of 6 pyrrolizidine alkaloids (PAs) have been isolated from 2 *Werneria* species. Alkaloids N-oxoretrosecine (**1**) and N-oxoretorsine (**4**) have been isolated from *W. decora* (Lock de Ugaz et al., 1990), while alkaloids retrorsine (**2**), senecionine (**3**), **4**, integerrimine (**5**) and N-oxorosmarinine (**6**) were reported for *W. nubigena* (Roeder et al., 1992; Piacente et al., 1997).


The presence of PAs is important from the chemotaxonomic point of view of *Werneria*, as it is generally regarded as a member of the tribe Senecioneae which is known to contain PAs. Pyrrolizidine alkaloids contain an azabicyclo [3,3,0] octane as a basic structure; they generally are found as esters with rare mono- or di-basic acids, the necic acids, producing a 12-membered macrocycle (**2-6**). It is also worth noting that, in general, the PAs accumulate in the plant as polar *N*-oxides (**1,4,6**).


### Benzofurans

Benzofurans are commonly found in Asteraceae family. Benzofurans dihydroeuparin or 6-hydroxytremetone (**7**) (Lock de Ugaz et al., 1984; Palacios et al., 2018), 2,5-diacetyl-6-hydroxybenzofurane (**8**) (Lock de Ugaz & Peralta, 1988), toxol (**9**) (Piacente et al., 1992), toxolacetate (**10**) (Córdova et al., 1998; de Marchese et al., 2007), toxolangelate


(11) (de Marchese *et al.*, 2007) and lactone 2,3-dihydro-2-oxo-5-formyl-6-methoxy-7-(3'-methyl-3'-butenyl)-benzo-furane (12) (Castro *et al.*, 2000) have a methylketone group in *para* position to the oxygen of the heterocycle ring, while 2-acetyl-5-(2'-

hydroxyethyl)-6-hydroxybenzofurane (13) (Castro *et al.*, 2000) has another substituent. These benzofurans were found in *W. ciliolata*, *W. poposa*, *W. nubigena* and *X. poposum*.







(7)




(8)



- (9) R = OH
- (10) R = acetyl
- (11) R = angelicyl



(12)








(13)

**Benzopyran derivatives (Chromenes)**

The basic structure of chromenes, which are benzopyrans derivatives, is a joined benzene and pyrane rings with two dimethyl group in position 2 and the methyl ketone group in *para* position to the oxygen of the heterocycle ring, as observed in 2,2-dimethyl-6-acetyl-8-(3'-hydroxy-3'-methyl-*trans*-but-1'-enyl)-chroman-4-one (**14**), 2,2-dimethyl-6-acetyl-8-(3'-hydroxy-3'-methyl-*trans*-but-1'-enyl)-chrom-3-ene (**15**) and 2,2-dimethyl-6-acetyl-3-chromene (**16**).

Sometimes this substituent is different like in 9-E-werneria chromene (**17**), 9-Z-werneria chromene (**18**), 8-[3',3'-dimethylallyl]-werneria chromene (**19**) and 8-[3',3'-dimethyl-3'-hydroxyallyl]-werneria chromene (**20**).





Compounds **14** and **15** were found in *W. nubigena* (Piacente *et al.*, 1997); **16** in *X. incisum* (de Marchese *et al.*, 2007) and **17-20** in *W. stuebelli* (Bohlmann *et al.*, 1984). Benzopyran **14** was reported as a new structure by Piacente *et al.* (1997).


**(14)****(15)****(16)****(17)** 9,10-E**(18)** 9,10-Z**(19)** R = 3', 3' -dimethylallyl**(20)** R = 3', 3'-dimethyl-3-hidroxyallyl

***p*-Coumarates and derivatives**


*p*-coumaric acid is formed through the deamination of L-tyrosine, one simple derivative is the compound methyl *p*-coumarate (**21**). Other *p*-coumarates like methyl 3-[3', 3'-dimethyl-3'-hydroxyallyl]-*p*-coumarate-*O*-methyl ether (**22**), ethyl-3-senecioid-*p*-coumarate (**23**), methyl-3-[3'-methyl-1'-hydroxy-but-

3'-en-1'-yl]-*p*-coumarate (**24**), Plicatin A (**25**) and Plicatin B (**26**) contain substituents in C-3 of the aromatic ring. Compounds **22-24** were reported for *W. stuebelli* (Bohlmann *et al.*, 1984), while Plicatin A and B were isolated from *W. dactylopylla* (Bravo *et al.*, 2009).





(25)




(26)

### Coumarins


Coumarins are phenolic compounds that are much extended in the plant kingdom; however, only five coumarins were reported so far for *Werneria* species: aesculetin (27), aesculetin-7-*O*- $\beta$ -D-glucopyranoside (28), scopoletin (29), 6-hydroxy-7-methoxycoumarin (30) and 7-hydroxy-8-methoxycoumarin (31). These

coumarins were found in *W. ciliolata*, *W. dactylophylla*, *W. poposa* and *W. staffordiae* (Bonilla *et al.*, 1991; Ponce & Gros, 1991; Piacente *et al.*, 1994; Chavez & Lock de Ugaz, 1997; Córdova *et al.*, 1998).



(27) R = OH

(28) R = Glc


(29) R<sub>1</sub> = CH<sub>3</sub>O, R<sub>2</sub> = OH, R<sub>3</sub> = H(30) R<sub>1</sub> = OH, R<sub>2</sub> = CH<sub>3</sub>O, R<sub>3</sub> = H(31) R<sub>1</sub> = H, R<sub>2</sub> = OH, R<sub>3</sub> = CH<sub>3</sub>O

### Flavonoids

Flavonoids are one of the most numerous and widely distributed compounds in plants which exhibit diverse biological activities. They appear usually as mixtures of aglycones and glycosides containing different classes and number of sugar units. The flavonoids reported for *Werneria* species were quercetin (**32**), rutin (**33**), Quercetin-3-*O*-glucorhamnoside (**34**), Quercetin-7-*O*-rhamnoside (**35**), Quercetin 3-*O*-( $\beta$ -D

glucopiranosyl-1'' $\rightarrow$ 2'')- $\beta$ -D-galactopyranoside (**36**) and hesperidin (**37**).

Compounds **32** and **33** were isolated from *W. poposa* and *W. dactylophylla* (Bonilla et al., 1991; Córdova et al., 1998), **34** and **36** from *W. staffordiae* (Chavez & Lock de Ugaz, 1997), while compounds **35** and **37** were reported in *W. dactylophylla* (Bonilla et al., 1991).




(32) R = R<sub>1</sub> = H

(33) R = H, R<sub>1</sub> = Rha-Glc

(34) R = H, R<sub>1</sub> = Glc-Rha

(35) R = Rha, R<sub>1</sub> = H

(36) R = OH, R<sub>1</sub> = Glc-Gal




(37)

### *p*-hydroxyacetophenones

*p*-hydroxyacetophenone (**38**) and its glucopyranoside (**39**) were isolated from *W. nubigena* and *W. poposa* (Piacente et al., 1997; Córdova et al., 1998). Compounds 3-(2'-hydroxy-isopent-3'-enyl)-4-hydroxy-acetophenone (**40**) and 3-(2',3'-dihydroxy-


isopentyl)-4-hydroxy-acetophenone (**41**) were reported in *W. ciliolata* (Piacente et al., 1992); while 4-hydroxy-3-(isopenten-2'-yl)-acetophenone (**42**) and 4-hydroxy-3-methoxy-acetophenone (**43**) were found in *X. incisum* and *X. poposum* (de Marchese et al., 2007; Palacios et al., 2018).




(38) R = R<sub>1</sub> = H

(39) R = Glc, R<sub>1</sub> = H

(43) R = H, R<sub>1</sub> = OCH<sub>3</sub>




(40)



(41)






(42)

### Caffeoylquinic acids

Caffeic acid is formed after a *p*-coumaric acid oxidation which introduces an OH group in *ortho*-position to the existent OH. They usually are alone as caffeic acids or combined with quinic acid to form *O*-

caffeoylquinic acids (44) or di-*O*-caffeoylquinic acids (45); in general they are called phenolcarboxylic acids and are very common in plants, in *Werneria* genus they have been reported in *W. nubigena* (Piacente *et al.*, 1997).











(44) R = H

(45) R = 3-*O*-caffeoylquinic acid

### Volatile terpenoids

The essential oils of *X. poposum* (= *W. poposa*) from Argentina and Chile have been studied by GC-MS by González *et al.* (2003, 2012), Mercado *et al.* (2015), Benites *et al.* (2011) and Abella *et al.* (2000). A sample of *X. incisum* from Argentina was studied by de Marchese *et al.* (2007). It is clear that we cannot compare these results because there are many factors that can influence the composition of the essential oils; i.e. vegetative cycle, environmental factors, extraction

Methods, among others. In general, the essential oils contain 20-60 compounds; we are reporting here only those with a yield greater than 5%. These volatile compounds include terpenoids that are found frequently in other essential oils like  $\beta$ -pinene (46),  $\beta$ -citronellol (47),  $\delta$ -cadinene (48),  $\gamma$ -cadinene (49), T-cadinol (50),  $\alpha$ -cadinol (51); as well as terpenoids that are less common like the tricyclic kessane (52), the alcohol incisol (53) and the macrocyclic (11C ring) compound zerumbone (54). Dihydroeuparin (7) is sometimes obtained as part of the essential oil.



### Diterpenes


Diterpenes **55-70** found in *Werneria* genus correspond mainly to *ent*-kauran and *ent*-manoyl oxide patterns. The prefix *ent* is used to indicate enantiomeric. Some new structures were found in *W. ciliolata* (Piacente *et al.*, 1994): 16 $\alpha$ ,17-dihydroxy-*ent*-kauran-19-al (**56**) and 16 $\alpha$ -methoxy-17-hydroxy-*ent*-kauran-19-al (**57**), 17-nor-16-oxo-kauran-19-oic acid (**59**), *ent*-3 $\beta$ ,16-dihydroxy-13-*epi*-manoyl oxide (**67**), 14,15-dihydroxy-*ent*-manoyloxide and acid 17-al-*ent*-kauran-19-oic acid (**70**), this last compound is a dimeric

structure between a kauranoic acid and manoyl oxide units linked by an oxygen atom.

Other known kauran derivative structures found were: kauran-16 $\alpha$ -ol (**55**) from *W. poposa* (Córdova *et al.*, 1998), **55** and (-)-kaur-16-en-19-oic acid (**58**) from *W. decora* (Lock de Ugaz *et al.*, 1990), 17-hydroxy-16 $\alpha$ -*ent*-kauran-19-oic acid (**61**) from *W. staffordiae* (Chavez & Lock de Ugaz, 1997), *ent*-kaur-16(17)-en-19-oic acid (**60**), 17-hydroxy-*ent*-kaur-15(16)-en-19-oic acid (**62**), 17-hydroxy-15 $\alpha$ , 16 $\alpha$ -epoxy-*ent*-kauran-19-oic-acid (**63**) and 17-al-*ent*-kaur-

15(16)-*ent*-19-oic-acid (**64**) from *W. ciliolata* (Piacente *et al.*, 1994). Manoyloxy compounds like *ent*-13-*epi*-manoyl oxide (**65**), *ent*-16-hydroxy-13-*epi*-manoyl oxide (**66**), *ent*-14 $\xi$ ,15 $\xi$ -epoxy-13-*epi*-manoyl


oxide (**68**) and *ent*-16-hydroxy-14 $\xi$ ,15 $\xi$ -epoxy-13-*epi*-manoyl oxide (**69**) were found in *Werneria dactylophylla* (De Tommasi *et al.*, 1992). Compound **68** was isolated for the first time from a natural source.



(**55**)  $R_1 = R_3 = \text{CH}_3$ ,  $R_2 = \text{OH}$

(**56**)  $R_1 = \text{CHO}$ ,  $R_2 = \text{OH}$ ,  $R_3 = \text{CH}_2\text{OH}$


(**57**)  $R_1 = \text{CHO}$ ,  $R_2 = \text{OCH}_3$ ,  $R_3 = \text{CH}_2\text{OH}$



(**58**)  $R_1 = \text{COOH}$ ,  $R_2 = \text{CH}_3$ ,  $\text{C}_{16}\text{-R}_3 = \text{=}$

(**59**)  $R_1 = \text{CH}_3$ ,  $R_2 = \text{COOH}$ ,  $\text{C}_{16}\text{-R}_3 = \text{c=O}$

(**60**)  $R_1 = \text{CH}_3$ ,  $R_2 = \text{COOH}$ ,  $\text{C}_{16}\text{-R}_3 = \text{=}$




(**61**)  $R_1 = \text{CH}_2\text{OH}$ ,  $R_2 = \text{H}$

(**62**)  $R_1 = \text{CH}_2\text{OH}$ ,  $\text{C}_{15}\text{-C}_{16} = \text{=}$

(**63**)  $R_1 = \text{CH}_2\text{OH}$ ,  $\text{C}_{15}\text{-C}_{16} = \text{}$

(**64**)  $R_1 = \text{CHO}$ ,  $\text{C}_{15}\text{-C}_{16} = \text{=}$




(**65**)  $R_1 = \text{H}$ ,  $R_2 = \text{CH}_3$

(**66**)  $R_1 = \text{H}$ ,  $R_2 = \text{CH}_2\text{OH}$

(**67**)  $R_1 = \text{OH}$ ,  $R_2 = \text{CH}_2\text{OH}$

(**68**)  $R_1 = \text{H}$ ,  $R_2 = \text{CH}_3$ ,  $\text{C}_{14}\text{-C}_{15} = \text{}$

(**69**)  $R_1 = \text{H}$ ,  $R_2 = \text{CH}_2\text{OH}$ ,  $\text{C}_{14}\text{-C}_{15} = \text{}$







### Triterpenes and Steroids

Few triterpenes (**71- 74**) and steroids have been detected or isolated from *Werneria* species. Triterpenes are found in many different skeletons mainly as tetracyclic and pentacyclic rings. Very common pentacyclic triterpenes like  $\beta$ -amirin (**71**), lup-20(29)-en-3-one (**72**), friedours-7-en-3-one (**73**), friedolean-6-en-3-one (**74**) were reported for *W.*

*nubigena* (Castro *et al.*, 2000).  $\beta$ -amirin was also isolated from *W. pigmea* (Aguilar *et al.*, 2000).

In these same species, only  $\beta$ -sitosterol and stigmaterol derivatives like stigmast-5,23-dien-3 $\beta$ -ol, stigmast-5,28-dien-3 $\beta$ -ol, stigmast-5,22-dien-3-ol and stigmast-5-en-3-one were reported (Castro *et al.*, 2000; Aguilar *et al.*, 2000).






### Hydrocarbons and acids

We considered not as important to describe the hydrocarbons and carboxylic acid open chains that have been detected or isolated in *Werneria* or

*Xenophyllum* species, except for isatinecic acid (75) (Lock de Ugaz *et al.*, 1990), a dicarboxylic acid which can form a macrocyclic diester (12C) with the alkaloid retronecine and other derivatives.



### Biological activities

A whole plant decoction of *Werneria villosa* is traditionally used for improving blood circulation, to treat general body pain and uterus swelling (Macía *et al.*, 2005).

*Werneria dactylophylla* is a plant used in Peruvian traditional medicine as a home remedy for inflammatory and gastrointestinal diseases (Valdizan

& Maldonado, 1922).

*Werneria poposa* is used mainly as an infusion of leaves and stems for the treatment of “soroche” (mountain sickness). Other uses include as a digestive or to treat coughs and bronchitis (Abella *et al.*, 2000). Martínez *et al.* (2006) reported that *W. incisa* and *W. poposa* are used in traditional Chilean medicine for the treatment of hypertension and mountain sickness.

*Werneria nubigena* is used as a tea because of its digestive and antihelmintic properties (Fournet *et al.*, 1994). The ethanol extract of the whole plant of *Werneria nubigena* showed activity *in vitro* against *Leishmania braziliensis*, *L. amazonensis* and *L. donovani* at a concentration of 100 µg/mL (Fournet *et al.*, 1994; Rocha *et al.*, 2005).

Erazo *et al.* (2006) found that, at a concentration of 100 µg/mL, the essential oil, hexane, dichloromethane and methanol extracts of *Xenophyllum poposum* showed antimicrobial activity *in vitro* against *Staphylococcus aureus*, *Bacillus subtilis* and *Saccharomyces cerevisiae*. The topical application of dichloromethane and hexane extracts of *X. poposum* decreased 12-*O*-tetradecanoylphorbol 13-acetate (TPA)-induced mouse ear edema by 80.9% and 60.6%, respectively. These results were compared to indomethacin, which reduced edema by 92.9%. These extracts showed also good analgesic activity in the acetic acid-induced writhing test. They caused an inhibition in the writhing response by 62.2% and 57.1%, respectively, which were similar to the inhibition showed by ibuprofen (76.5%).

The hydroalcoholic extract of *X. poposum* administered orally at 40 mg/kg significantly reduced blood pressure and heart rate in normotensive rats. The hypotensive activity is due in part to the decrease in heart rate and increased vasodilation. The compounds that caused the greatest vasodilation were 4-hydroxy-3-(3-methyl-2-butenyl) acetophenone (**42**) and dihydroeuparin (**7**). These results could support the use *X. poposum* in traditional medicine in the treatment of altitude sickness and hypertension (Palacios *et al.*, 2018).

The essential oil of *X. poposum* collected in Argentina showed good antimicrobial activity *in vitro* against the bacterium methicillin-resistant *Staphylococcus aureus* and the fungi *Trichophyton rubrum* and *Aspergillus fumigatus*, the MIC values were 0.25, 0.05 and 0.025 mg/mL, respectively. The antimicrobial activity of the methanol extract was very similar to the one reported for the essential oil. The most active compound isolated from *X. poposum* methanol extract was dihydroeuparin (**7**), this compound was very active against *A. fumigatus*, although its MIC value was equal to that of the essential oil (0.025 mg/mL) (González *et al.*, 2012).

Piacente *et al.* (2004) tested the anti-HIV activity *in vitro* of the compounds isolated from *Werneria ciliolata* and *W. dactylophylla*. The *p*-hydroxyacetophenones and benzofurans were inactive,

however, interesting results were obtained with the compound dihydroeuparin (**7**) that was isolated from *W. dactylophylla* (Piacente *et al.*, 1994). At a concentration of 16 µg/mL, dihydroeuparin was able to reduce by 50% the P24 antigen in infected cell cultures, while at a concentration of 40 µg/mL it inhibited by 50% the growth of the uninfected C8166 cells (TC<sub>50</sub>). The drug AZT was used as a positive control; this compound was very active *in vitro* (EC<sub>50</sub>= 0.016 µg/mL) and showed very low toxicity (TC<sub>50</sub>>1000 µg/mL) (Piacente *et al.*, 1994).

*Artemia salina*, the brine shrimp larva, was used to determine the toxicities of Plicatin A and Plicatin B. The medium lethal concentrations (LC<sub>50</sub>) of both compounds were 0.96 ± 0.17 ppm (Bravo *et al.*, 2009).

Ávila Illanes *et al.* (2011) found a LD<sub>50</sub> greater than 5000 mg/kg for the aqueous extract of *Werneria dactylophylla* (pupusa) when administered orally to albino Swiss mice.

Although *Werneria* and *Xenophyllum* extracts are used in traditional medicine and show no toxicity in animals, it has to be taken into account that they contain Pyrrolizidine alkaloids (PAs) that can cause veno-occlusive disease and liver damage. The acute toxic effects of PAs are characterized by hemorrhagic necrosis, hepatomegaly and ascites; while chronic intoxication causes necrosis, cirrhosis, liver failure and death. PAs with 1,2 double bond, like for example retronecine and **1**, are associated with hepatic toxic effects. PAs with cyclic diesters, like in structures **2-6**, are considered the ones with the highest toxicity (Moreira *et al.*, 2018).

The compound 4-hydroxy-3-(isopenten-2'-yl)-acetophenone (**42**) is the main secondary metabolite of the aerial parts of *Senecio nutans* and *X. poposum*. At a concentration of 5 mg/L this compound showed a synergistic effect in combination with Fluconazole or with Caspofungin against *Candida albicans* ATCC 10231. Such synergistic activity could be explained by the fact that these compounds have different mechanisms of action. Fluconazole inhibits the synthesis of ergosterol, Caspofungin inhibits cell wall β-glucan synthesis and 4HMBA inhibits *C. albicans* filamentation (Soberón *et al.*, 2015).

The benzofuran derivative dihydroeuparin can be found in *Werneria ciliolata* and in the shrub *Senecio graveolens* with isolation yields of near 3%. Photochemical and photophysical results indicate that this benzofuran can be considered as a sunscreen for the 270-340 nm zone, with high photostability, low

water solubility and high solubility in miscellar system (Ortega et al., 2000).

## CONCLUSIONS

*Werneria* and *Xenophyllum* species are plants that are widely used in traditional medicine for the treatment of hypertension, mountain sickness and inflammatory disorders. Some interesting anti-HIV, antifungal and photoprotective activities have been reported. However, it has to be noted that these plants contain pyrrolozidine alkaloids that could be hepatotoxic. Only 9 species of *Werneria* and 2 of *Xenophyllum* have been studied for their chemical and biological activities; thus, there are more than 50 species that deserve to be further investigated.

## REFERENCES

- Abella L, Cortella AR, Velasco-Negueruela A, Pérez-Alonso MJ. 2000. Ethnobotany, volatile oils and secretion tissues of *Werneria poposa* from Argentina. **Pharm Biol** 38: 197 - 203.
- Aguilar R, Rodríguez F, García M. 2000. **Identificación de metabolitos secundarios de *Werneria pygmaea* Gilles por CG-EM.** Libro de Resúmenes del XXIV Congreso Latinoamericano de Química. Lima, Perú.
- Avila Illanes JA, Ruiz Pinell G, Tórrez Choque C, Mollinedo Portugal P, Vila Castro JL, Bugiardini M, Sauvain M, Gonzáles E, Bravo JA. 2011. Evaluación de la toxicidad aguda a dosis fijas del extracto de *Werneria dactylophylla* (Pupusa). **Rev Bol Quím** 28: 87 - 90.
- Beltrán H. 2016. Sinopsis del género *Xenophyllum* (Asteraceae: Senecioneae) del Perú. **Arnaldoa** 23: 351 - 362.
- Beltrán H. 2017. Sinopsis del género *Werneria* (Asteraceae: Senecioneae) del Perú. **Arnaldoa** 24: 45 - 62.
- Beltrán H, Leiva S. 2018. *Werneria microphylla* (Asteraceae, Senecioneae), a new species from the Andean marshes of Peru. **Phytotaxa** 372: 293 - 300.
- Benites J, Lopez J, Rojo LE, Diaz P, Rojas M, Fuentes R, Moiteiro C, Venancio F. 2011. Chemical composition of the essential oil of *Xenopyllum poposum*. **Chem Nat Compd** 46: 988 - 989.
- Bohlmann F, Zdero C, King RM, Robinson H. 1984. Prenylated *p*-coumarates from *Werneria stuebelli*. **Phytochem** 23: 1135 - 1137.
- Bonilla PE, Lock de Ugaz O, Jurupe H. 1991. Estudio químico y biológico de la *Werneria dactylophylla*. **Bol Soc Quím Peru** 57: 182 - 188.
- Bravo JA, Camacho V, Rendón WJ. 2009. Chemical education: Structural elucidation Plicatin A and B from *Werneria dactylophylla*. **Rev Bol Quím** 26: 63 - 76.
- Castro R, García M, Rodríguez F. 2000. **Identificación de metabolitos secundarios en la *Werneria nubigena* por CG-EM.** Libro de Resúmenes del XXIV Congreso Latinoamericano de Química. Lima, Perú.
- Chávez R, Lock de Ugaz O. 1997. Contribución al estudio químico de la *Werneria staffordiae* Sandwith. Análisis por RMN de sus metabolitos secundarios. **Bol Soc Quím Peru** 63: 211 - 223.
- Córdova A, Lock O, Jurupe H. 1998. Estudio químico farmacológico de la *Werneria poposa* Philippi. **Bol Soc Quím Peru** 64: 264 - 272.
- de Marchese MJA, de Heluani CS, Catalan CAN, Griffin CA, Vaughn JB, Herz W. 2007. Incisol, an alcohol with a novel sesquiterpene skeleton from *Xenophyllum incisum*. **Biochem Syst Ecol** 35: 169 - 175.
- De Tommasi N, Aquino R, de Simone F, Piacente S, Pizza C. 1992. Diterpenes from *Werneria dactylophylla*. **Phytochemistry** 31: 1042 - 1044.
- Erazo S, Gogoux F, Delporte C, Backhouse N, Negrete R, Zaldívar M, Belmonte E, Cayunao C, Bachiller I, Bandoni A, Di Leo Lira P, van Baren C, Gianninoto H, García R. 2006. Actividades biológicas de *Xenophyllum poposum* Phil. (Asteraceae), planta del altiplano chileno. **Rev Fitoterapia** 6: 165 - 166.
- Fournet A, Barrios AA, Muñoz V. 1994. Leishmanicidal and trypanocidal activities of Bolivian medicinal plants. **J Ethnopharmacol** 41: 19 - 37.
- Funk VA. 1997a. *Xenophyllum*, a new Andean genus extracted from *Werneria* s.l. (Compositae: Senecioneae). **Novon** 7: 235 - 241.
- Funk VA. 1997b. *Misbrookea*, a new monotypic genus removed from *Werneria* s.l. (Compositae: Senecioneae). **Brittonia** 49: 110 - 117.
- González A, Poch M, Amani S, Tracanna M, Heluani C, Catalán C. 2003. **Mono y sesquiterpenoides del aceite esencial de *Xenophyllum poposum* (Phil.) V. A. Funk.** IV

- Encuentro Regional de Plantas Medicinales del NOA. Resúmenes en CD, No 22. Horco Molle, Tucumán, Argentina.
- González AM, Tracanna MI, Amani SM, Schuff C, Poch MJ, Bach H, Catalán CAN. 2012. Chemical composition, antimicrobial and antioxidant properties of the volatile oil and medicinal extract of *Xenophyllum poposum*. **Nat Prod Commun** 7: 1663 - 1666.
- Linares-Perea E, Campos-de la Cruz J, Galán de Mera A. 2014. *Anticona*, un nuevo género andino relacionado con *Werneria*. (Compositae). **Perspectiva** 15: 99 - 102.
- Lock de Ugaz O, Hajar A, Borges del Castillo J, Seligmann O, Wagner H. 1984. Dihydroeuparine from *Werneria ciliolata*. **Fitoterapia** 55: 248.
- Lock de Ugaz O, Peralta A. 1988. Benzofurano de la *Werneria ciliolata*. **Rev Lat Quim** 19: 71 - 72.
- Lock de Ugaz O, Franco J, Seminario G, Delle Monache F, Millan B, Ubillas RP, Schlemper EO, Tempesta MS. 1990. Alkaloids and diterpenoids from *Werneria decora*. **Phytochem** 29: 2373 - 2375.
- Lock Sing O. 2006. Diversidad química en el género *Werneria*. **Bol Soc Quim Perú** 72: 32-43.
- Macía MJ, García E, Vidaurre PJ. 2005. An ethnobotanical survey of medicinal plants commercialized in the markets of La Paz and El Alto, Bolivia. **J Ethnopharmacol** 97: 337 - 350.
- Martínez JL, Calvo CA, Laurido C. 2006. Medicinal plants used in Chile for the treatment of hypertension and mountain sickness. **Afr J Trad Compl Alt Med** 3: 50 - 58.
- Mercado MJ, Lizárraga E, Catalán CAN, Pomessa GI. 2015. Composición del aceite esencial de *Xenophyllum poposum* (Asteraceae) en la provincia de Catamarca y Tucumán. Estructuras foliares y caulinares asociadas a su síntesis. **Lilloa** 52: 56-61.
- Moreira R, Pereira DM, Valentão P, Andrade PB. 2018. Pyrrolizidine alkaloids: Chemistry, pharmacology, toxicology and food safety. **Int J Mol Sci** 19: 1668, 1 - 22.
- Ortega PA, Guzmán ME, Vera LR, Velásquez JA. 2000. Dihydroeuparin as a sunscreen. **Bol Soc Chil Quim** 45: 629 - 636.
- Palacios J, Quispe C, Paredes A, Cifuentes F. 2018. Uso potencial de *Xenophyllum poposum* (Phil) V.A Funk (Compositae) en el tratamiento del mal de altura e hipertensión. **Med Plant Commun** 1: 24.
- Piacente S, Aquino R, De Tommasi N, Lock de Ugaz O, Chavez Orellana H. 1992. *p*-hydroxyacetophenone derivatives from *Werneria ciliolata*. **Phytochemistry** 31: 2182 - 2184.
- Piacente S, Aquino R, De Tommasi N, Pizza C, Lock de Ugaz O, Chavez Orellana H, Mahmood N. 1994. Constituents of *Werneria ciliolata* and their *in vitro* anti-HIV activity. **Phytochem** 36: 991 - 996.
- Piacente S, Herrera N, De Simone F, Lock de Ugaz O, Pizza C. 1997. Benzopyran derivatives from *Werneria nubigena*. **Phytochem** 46: 795 - 797.
- Ponce MA, Gros E. 1991. Prenylated aromatic ketones from *Werneria poposa*. **Anal Asoc Quim Arg** 79: 197 - 200.
- Rocha LG, Almeida JRGS, Macêdo RO, Barbosa-Filho JM. 2005. A review of natural products with antileishmanial activity. **Phytomedicine** 12: 514 - 535.
- Roeder E, Bourauel T, Theisen I. 1992. Pyrrolizidine alkaloids from *Werneria nubigena*. **Nat Toxins** 1: 81 - 83.
- Soberón JR, Lizarraga EF, Sgariglia MA, Carrasco Juárez MB, Sampietro DA, Altabef AB, Catalán CAN, Vattuone MA. 2015. Antifungal activity of 4-hydroxy-3-(3-methyl-2-butenyl) acetophenone against *Candida albicans*: evidence for the antifungal mode of action. **Antonie van Leeuwenhoek** 108: 1047 - 1057.
- Valdizan H, Maldonado A. 1922. La Medicina Popular Peruana, Vol. II. Imprenta Torres Aguirre, Lima, Perú.